

# POISON

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS  
BEFORE OPENING  
OR USING



WEEDFORCE®

# Dart® 100 SC

GRUB SPIDER ANT  
INSECTICIDE



**ACTIVE CONSTITUENT: 100 g/L BIFENTHRIN**

**GROUP 3A INSECTICIDE**

For the control of a range of urban interior & exterior pests, including spiders, mosquitoes, flies, and other pests, for protection of structures from subterranean termite damage and for control of various pests in turf, orchards and ornamentals as specified in the directions for use table.

**IMPORTANT: READ THIS LEAFLET BEFORE USE**

WEEDFORCE Pty Ltd ACN 602 207 152   
2072 Logan Rd Mt Gravatt QLD 4122  
P: 0488 997 923 [www.weedforce.com.au](http://www.weedforce.com.au)

APVMA Approval No:  
92587 / 135972

 Registered Trademark of Weed Force Pty Ltd  
© Weed Force Pty Ltd 2022

## WEED FORCE DART 100SC DIRECTIONS FOR USE

### INDEX

RESTRAINTS.....	3
USE RATES FOR INTERNAL AND EXTERNAL AREAS, TURF/LAWNS .....	4-7
TABLE A: USE RATES FOR CONTROL OF SUBTERRANEAN TERMITES .....	8-9
TABLE B: CRITICAL COMMENTS FOR USE AGAINST SUBTERRANEAN TERMITES .....	10-11
DIRECTIONS FOR USE- ORCHARDS AND ORNAMENTALS .....	12-15
GENERAL INSTRUCTIONS .....	16
MIXING & APPLICATION.....	16
COMPATIBILITY.....	17
CRITICAL APPLICATION DETAILS -TERMITES .....	17
INSECTICIDE RESISTANCE WARNING .....	19
PRECAUTIONS AND RE-ENTRY PERIODS .....	20
PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT .....	20
PROTECTION OF PETS AND LIVESTOCK .....	20
STORAGE AND DISPOSAL.....	21
SAFETY DIRECTIONS.....	21
FIRST AID .....	21
SAFETY DATA SHEET .....	22
CONDITIONS OF SALE.....	22

### DIRECTIONS FOR USE:

#### RESTRAINTS:

DO NOT use this product at less than indicated label rates.

DO NOT apply to soils if excessively wet or immediately prior to or after heavy rain to avoid run-off of the chemical.

DO NOT apply to mud, sand, mangrove or aquatic habitat.

DO NOT apply as an Ultra Low Volume (ULV) or via thermal fogging treatment.

DO NOT use in situations where predatory mites are established and providing effective mite control.

DO NOT apply if rainfall is expected before spray deposits dry on leaf surfaces.

DO NOT use in cavity walls.

#### Orchards and Ornamentals

DO NOT use in situations or orchards where predatory mites are established and providing effective mite control.

DO NOT apply by aircraft.

DO NOT apply if rainfall is expected before spray deposits dry on leaf surfaces.

## Use rates for INTERNAL AND EXTERNAL AREAS, TURF/LAWNS

Situation	Pest	Rate	Critical Comments
Internal & external areas & surrounds of domestic, commercial, public & industrial buildings and structures	Spiders	25-50 mL/10 L	Use the higher rate in situations where pest pressure is high, when rapid knockdown and/or maximum residual protection is desired. Pay particular attention to protected dark areas such as cracks and crevices, under floors, eaves, and other known hiding or resting places. For overall band surface spray, apply as a coarse, low pressure surface spray to areas where spiders hide, frequent and rest. Spray to the point of run-off using around 5 L of spray mixture per 100 m <sup>2</sup> and ensuring thorough coverage of the treated surfaces. For crack and crevice treatment use an appropriate solid stream nozzle. For maximum spider control use a two part treatment. <ol style="list-style-type: none"> <li>1. Crack and crevice.</li> <li>2. Overall band spray of surfaces.</li> </ol>
	Papernest Wasps	50 mL/10 L	Apply prepared solution to the point of runoff directly to the papernest ensuring thorough and even coverage. When all adult wasps have been knocked-down the nest may be safely removed from the structure.
Internal & external areas & surrounds of domestic, commercial, public & industrial buildings and structures  - suitable for residual surface treatments	Ants (excluding Red Imported Fire Ants), Cockroaches, Mosquitoes, Fleas, Flies, Ticks (excluding the paralysis tick <i>Ixodes holocyclus</i> ) (Adults & Nymphs)	50-100 mL/10 L	Use the higher rate in situations where pest pressure is high, when rapid knockdown and/or maximum residual protection is desired. The lower rate may be used for follow-up treatments. For indoor use, pay particular attention to protected dark areas such as cracks & crevices, behind or under sinks, stoves and refrigerators, furniture, pipes, comices, skirting boards and other known hiding or resting places. DO NOT use a surface spray. <b>On non-porous surfaces</b> apply as a coarse spray at the rate of 1L solution per 20 m <sup>2</sup> . When treating non-porous surfaces do not exceed the point of run-off. <b>On porous surfaces</b> use through power equipment, spray at the rate of 1 L of solution per 20 m <sup>2</sup> . When treating porous surfaces do not exceed the point of run-off.  <b>Ants:</b> To control ants apply to trails and nests. Repeat as necessary. <b>Fleas and ticks:</b> To control fleas and ticks, apply prepared solution to outside surfaces of buildings and surrounds including but not limited to foundations, verandas, window frames, eaves, patios, garages, pet housing, soil, turf, trunks or woody ornamentals or other areas where pests congregate or have been seen. <b>Flies and Mosquitoes:</b> To control flies and mosquitoes apply prepared solution to surfaces where insects rest or harbour. Repeat as necessary. <b>Perimeter treatments:</b> Apply the prepared solution to a band of soil or vegetation two to three meters wide around and adjacent to the structure. Also treat the foundation of the structure to a height of approximately one metre. Use a spray volume of 5 to 10 L per 100 m <sup>2</sup> . Higher volumes of water may be needed if organic matter is present or foliage is dense.

Use rates for **INTERNAL AND EXTERNAL AREAS, TURF/LAWNS** Continued

Situation	Pest	Rate	Critical Comments
Turf (eg lawns, commercial turf farms, parks, recreational areas, bowling greens, sports fields)	Lawn armyworm ( <i>Spodoptera maurita</i> ), Sod webworm ( <i>Herpetogramma licarsisalis</i> )	1.2 L/ha (12 mL/100 m <sup>2</sup> )	Mix WEED FORCE DART 100SC GRUB SPIDER ANT INSECTICIDE in water and apply evenly over the area to be treated using spray application equipment. Use a minimum water volume of at least 200 L/ha (2L/100 m <sup>2</sup> ). To ensure optimum control, irrigate the treated area with up to 4mm of water soon after application. Inspect treated areas for continuing activity. Reapply as required. Where a rate range is indicated use lower rates under lower insect pressure and higher rates under higher insect pressure. Apply after mowing to minimise loss of insecticide in clippings.
	Argentine stem weevil adults ( <i>Listronatus bonariensis</i> )	1.2 – 2.4 L/ha (12-24 mL /100 m <sup>2</sup> )	
	African black beetle adults ( <i>Heteronychus arator</i> )	2.4-3.6 L/ha (24-36 mL/100 m <sup>2</sup> )	
	Billbug adults ( <i>Sphenophorus brunnipinnis</i> )	1.2-2.4 L/ha (12 - 24 mL/100 m <sup>2</sup> )	
	Black ant, coastal brown ant, funnel ant, meat ant, sugar ant and stinging ant only	1.2-4.4 L/ha (12-44 mL/100 m <sup>2</sup> )	
Domestic, Public, Commercial & Industrial areas	Subterranean Termites	Refer to Table A	Refer to Table B.

**TABLE A: Use rates for control of SUBTERRANEAN TERMITES**

Situations	All areas SOUTH of the Tropic of Capricorn (except Tas.)		All areas NORTH of the Tropic of Capricorn	
	Rate	Expected Protection Period *	Rate	Expected Protection Period *
<b>Perimeter Barriers</b> For new and existing buildings	1 L/100 L	At least 10 years	1.5 L/100 L	Up to 5 years
	500 mL/100 L	10 years	1 L/100 L	Up to 4 years
	250 mL/100 L	3 years	750 mL/100 L	Up to 3 years
<b>Protection of Poles &amp; Fence Posts</b>	500 mL/100 L	10 years	500 mL/100 L	Up to 2 years
			1.5 L/100 L	Up to 5 years
			1 L/100 L	Up to 4 years
<b>Nest Eradication</b>	500 mL/100 L	Not applicable	750 mL/100 L	Up to 3 years
			500 mL/100 L	Not applicable
<p><b>*Several factors contribute to the estimated length of protection provided for each termite treatment. The actual protection period will depend on the termite hazard, climate, soil conditions and rate of termiticide used. The need for retreatment is to be determined as a result of at least an annual inspection, or more frequently in high risk areas, by a qualified licensed Pest Control Operator.</b></p>				

**TABLE B: CRITICAL COMMENTS for use against SUBTERRANEAN TERMITES**

Situations	Critical Comments
<b>Perimeter Barriers</b> for new and existing buildings **	<ul style="list-style-type: none"> <li>Perimeter barriers (both horizontal and vertical, external and where required, internal or sub-floor) are an essential part of termite protection and must be installed at the completion of the building. Perimeter barriers should be installed around slabs, piers, substructure walls and external penetration points.</li> <li>Apply with suitable application equipment to form a continuous chemical barrier (both vertical and horizontal) around the structure and to a depth reaching to 80 mm below the top of the footings, where appropriate. The formation of the barrier may require a combination of several application techniques, including soil trenching and/or rodding and open wand applications.</li> <li>Chemical barriers that have been disturbed by construction, excavation and/or landscaping activities will need to be reapplied to restore continuity of the barrier.</li> </ul>
<b>Protection of Service Poles and Fence Posts</b>	<ul style="list-style-type: none"> <li>Create a continuous termiticide barrier 450 mm deep and 150 mm wide around the pole or post by soil injection or rodding. For new poles and posts, treat backfill and the bottom of the hole. Use 100 L of solution per m<sup>3</sup> of soil.</li> <li>Regular inspections should be undertaken to determine when and if retreatment is necessary. If disturbance of the barrier has occurred, retreatment of the area affected will be required.</li> <li>Posts and poles may also be drilled and injected with spray solution.</li> <li>Note: For existing poles and posts, it is impractical to treat the full depth and underneath of such poles and posts and therefore the possibility of future termite attack from below the treated area cannot be ruled out.</li> <li>For establishing trees create a continuous barrier totally encompassing the root ball of the establishing tree. Application may be made prior to planting by applying solution to pre-dug hole or after planting via soil rodding. Roots projecting out of the treated zone may be susceptible to termite attack and may provide entry into the tree without termites contacting treated soil.</li> <li>WEED FORCE DART 100SC GRUB SPIDER ANT INSECTICIDE is a non-systemic insecticide. DO NOT treat mature trees as it is impossible to provide a complete and continuous barrier under and around all tree roots.</li> </ul>

Situations	Critical Comments
<b>Eradication of Termite Nest</b>	<ul style="list-style-type: none"> <li>Locate nest and flood with insecticide solution. Trees, poles, posts and stumps containing nests may require drilling prior to treatment with termiticide solution. The purpose of drilling is to ensure the termiticide solution is distributed throughout the entire nest. Drill holes in live trees should be sealed with an appropriate caulking compound after injection.</li> </ul>
<b>Notes to Critical Comments</b>	
** Chemical barriers that have been disturbed by construction, excavation and /or landscaping activities will need to be reapplied to restore continuity of the barrier.	

**Note: The termiticide barrier provided by this product has a finite life. This together with the recommendation to undertake annual inspections must be stated on the durable notice required by the BCA, B1.3(j)(ii).**

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**

**DIRECTIONS FOR USE- ORCHARDS AND ORNAMENTALS**

CROP	PEST	STATE	RATE	CRITICAL COMMENTS
Apples	Apple Dimpling Bug ( <i>Campylomma liebknecht</i> ), Plague Thrips ( <i>Thrips imaginis</i> )	Qld, NSW, Vic, SA, WA only	8 mL or 16 mL/100 L	Apply when pest numbers reach accepted threshold levels. Applications should be made as early as possible during the blossoming period and early in the morning when bees are not actively foraging. Use the high rate for both knockdown and residual control. Only one application at this rate should be required per season. In orchards where appropriate crop monitoring facilities are available, the low rate may be used for knockdown control only. When this low rate is used, a second application at the low rate may be required to control re-infestation. Spray to run-off using a total spray volume of 1000 to 2500 L/ha depending on tree size.
Bananas	Banana Scab Moth ( <i>Nacoleia octasema</i> )	Qld only	200 mL/100 L	Apply 40 mL of prepared spray to each banana bell. Use a suitable bell injection instrument to inject the required volume directly into the bell as it emerges from the throat of the banana plant while in the upright position. The correct site for injection is in the top half to one third of the bell just below the distinct swelling where the male flower mass ends and the female flower cavity (bottom hand of fruit) start. Keep injection equipment clean and use lubricants sparingly. Monitor for thrips activity and treat only when thrips are active.
	Flower Thrips ( <i>Thrips florum</i> )	Qld, NSW only		
Pears	Two Spotted Mite ( <i>Tetranychus urticae</i> ), Pear Looper, Longtailed Mealy Bug Crawlers, Lightbrown Apple Moth, Codling Moth	Vic only	40 mL/100 L	Monitor the mite population from mid December onwards. Apply the product before the mite population reaches economic damage levels (ie around 20-30 motiles/25 leaves). A follow-up treatment may be required 3 to 4 weeks later. If more than 2 miticide applications are required use an alternative rotational miticide. Spray to run-off using a total spray volume of 2000-4000 L/ha depending on the tree size. NOTE: When using WEED FORCE DART 100SC GRUB SPIDER ANT INSECTICIDE in pears, it is not necessary to tank-mix additional insecticides for control of Codling Moth and Lightbrown Apple Moth, Pear Looper and Longtailed Mealy Bug Crawlers.

CROP	PEST	STATE	RATE	CRITICAL COMMENTS
Roses, Carnations & other ornamental plants	Two Spotted Mite ( <i>Tetranychus urticae</i> )	All States	28 or 40 mL/100 L	Apply at first sign of pest infestation and before pest populations build up to damaging levels. Repeat as necessary on a 10-14 day interval. Best results are obtained from preventative rather than curative applications. Where indicated, use the higher dosage for knockdown of established pest infestations or when longer residual activity is required. Spray to run-off using a spray volume of 1000-1500 litres per ha (10-15 L/m <sup>2</sup> ) covering both leaf surfaces.
	Aphids		20 mL/100 L	
	Caterpillars and Loopers including <i>Heliothis</i> (Corn Ear Worm, Native Budworm), <i>Helicoverpa</i> spp., Light Brown Apple Moth, <i>Epiphyas postvittana</i> , Geranium Plume Moth ( <i>Sphenarches anisodactylus</i> )		20 mL/100 L	Apply at first sign of pest infestation and before pest populations build up to damaging levels. Repeat as necessary on a 10-14 day interval. Best results are obtained from preventative rather than curative applications. Spray to runoff using a spray volume of 10-15 litres per 100 square metres covering both leaf surfaces.
	Whitefly ( <i>Trialeurodes vaporariorum</i> ), Poinsettia White Fly ( <i>Bemisia tabaci</i> Biotype B)		20-80 mL/100 L	Apply at first sign of pest activity and repeat at intervals of 7-10 days while pest pressure persists. More than three sprays may be required to control an existing infestation. Spray to run-off covering both leaf surfaces. Use the higher rate when pest pressure is high, when conditions favour pest development or when increased residual protection is required.
	Mealy Bug ( <i>Pseudococcus longispinus</i> )		20 mL/100 L	Apply at first sign of pest activity and repeat at intervals of 7-10 days while pest pressure persists. Spray to run-off covering both leaf surfaces.
	Plague Thrips ( <i>Thrips imaginis</i> , <i>Thrips simplex</i> and <i>Thrips hawaiiensis</i> )		20 mL/100 L	Apply at first sign of pest activity and repeat at intervals of 7-10 days while pest pressure persists. Ensure that flowers and buds sprayed. Spray to run-off. When buds are opening rapidly and pest pressure is high reducing the spray interval to 3-4 days will give better results. Monitor the population by regular inspection.
	Cutworm ( <i>Agrotis</i> spp.) in beds, containers and pots		1.2 L/ha 12 mL/100 m <sup>2</sup>	Spray evenly over the area to be treated. After application apply approximately 5 mm of sprinkler irrigation.
			20 mL/ 100 L	Apply as a drench at the rate of 2 litres of prepared spray per square metre of pot area.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**

**WITHOLDING PERIODS:**

**PEARS: DO NOT APPLY LATER THAN 14 DAYS BEFORE HARVEST**



## GENERAL INSTRUCTIONS

**Urban Pest Control:** Weed Force Dart 100SC Grub Spider Ant Insecticide has powerful knockdown and residual activity against various insect pests. Ants, cockroaches, fleas, flies, mosquitoes, spiders, ticks and wasps are controlled by direct contact with spray and also by residual action as they come in to contact with the treated surfaces.

**Termites:** The use of Weed Force Dart 100SC Grub Spider Ant Insecticide will help prevent and control subterranean termite infestations in and around buildings and structures when used in accordance with the Australian Standard AS 3660 Series, Termite Management. A dilute termiticide solution must be adequately dispersed into the soil to establish a barrier between the building and subterranean termites in the soil. The purpose of a termite barrier is to prevent concealed termite entry into the building.

The biology and behaviour of the termite species involved should be considered by the Pest Control Operator in determining which control measures are most appropriate to control and prevent termite infestation.

Best results are obtained when WEED FORCE DART 100SC GRUB SPIDER ANT INSECTICIDE is applied before pest populations build up to damaging levels.

**Turf:** This product can be used as a protective treatment when applied at regular intervals or as a knockdown treatment to control existing pests. Best results are obtained when the product is applied before pest populations build up to damaging levels.

### Mixing & Application

WEED FORCE DART 100SC GRUB SPIDER ANT INSECTICIDE is a suspension concentrate requiring dilution with water prior to use. Add the required quantity of WEED FORCE DART 100SC GRUB SPIDER ANT INSECTICIDE to water in the spray tank and mix thoroughly. Maintain agitation during both mixing and application. Ensure that spray lines are clean and free from contaminant pesticide, eg. herbicides.

**For Termites:** To facilitate even application of the termiticide solution over the area to be treated, the addition of a marker dye at label rates is recommended. On hard to wet soils, the penetration of the termiticide solution may be improved by the addition of a soil surfactant at label rates.

**For General Pest Control:** Application of WEED FORCE DART 100SC GRUB SPIDER ANT INSECTICIDE should be made with equipment calibrated to deliver a fine dilute spray in a suitable volume of water to ensure thorough coverage. Use suitable application equipment and preferably cone nozzle combinations to deliver the appropriate spray volume and FINE droplets. DO NOT apply as a fog or mist.

**For Vector Control Applications:** WEED FORCE DART 100SC GRUB SPIDER ANT INSECTICIDE can be applied by a variety of application techniques, such as power sprayers, hand held pump sprayers or backpack sprayers, which are operated to produce large droplets to minimise spray drift. However, the product must be not applied by ULV or fogging equipment, which are operated to produce small droplets that may result in off-target drift.

**For Turf Applications:** Application should be made with equipment calibrated to deliver a fine dilute spray in a suitable volume of water to ensure thorough coverage. Use suitable application equipment and preferably cone nozzle combinations to deliver appropriate spray volume and FINE droplets. DO NOT apply as a fog or mist. Maintain agitation during both mixing and application.

### COMPATIBILITY:

Weed Force Dart 100SC Grub Spider Ant Insecticide is compatible with commonly used fungicides such as Mancozeb, Antracol, Clorothalonil 500 and Copper (as Cupic Oxide)

### Surfactants

Weed Force Dart 100SC Grub Spider Ant Insecticide contains a surfactant. Additional surfactant may only be necessary on hard to wet plants and in high volume sprayers.

## CRITICAL APPLICATION DETAILS -TERMITES

### Treatment of Structures:

The pest control operator must notify site supervisor if any, and workers who come in contact with uncovered treated soil prior to laying the moisture membrane, to wear appropriate personal protective equipment, and to observe re-entry requirements. (For personal protective equipment refer to "SAFETY DIRECTIONS", and for re-entry refer to "PRECAUTIONS AND RE-ENTRY PERIODS", below).

The use of WEED FORCE DART 100SC GRUB SPIDER ANT INSECTICIDE will help prevent and control subterranean termite infestations in and around buildings and structures when used in accordance with the Australian Standard AS 3660 Series. Termite Management - A dilute termiticide solution must be adequately dispersed into the soil to establish a barrier between the building and subterranean termites in the soil. The purpose of a termite barrier is to prevent concealed termite entry into the building. The biology and behaviour of the termite species involved should be considered by the pest control operator in determining which control measures are most appropriate to control and prevent termite infestation. The application of WEED FORCE DART 100SC GRUB SPIDER ANT INSECTICIDE to form both horizontal and vertical chemical barriers must be in accordance with the Australian Standard AS 3660 Series. For treatment of new and existing buildings, both horizontal and vertical

barriers may be required, around and under the building. External perimeter barriers and where required, internal perimeter barriers are an essential part of this treatment. The purpose of a chemical termite soil barrier is to provide a continuous, no gap barrier between the building and the termite colony. It is therefore essential that the Pest Control Operator is familiar with the construction details of the building. For further details, refer to the "Horizontal Barrier Treatments" and "Vertical Barrier Treatments" statements in this leaflet and to the Australian Standard AS 3660 Series.

**Horizontal Barrier Treatments:** Use 5 L of solution per m<sup>2</sup> of soil. Apply the termiticide solution evenly to the soil surface area to ensure the provision of a continuous barrier with no gaps. To minimise drift, use low pressure, high volume spray equipment delivering large coarse droplets. On impervious soils where the application of 5 L/m<sup>2</sup> would cause excessive run-off, the application volume may be reduced provided the concentration of the solution is increased by a corresponding amount. For example, the volume of applied concentrate must remain constant at 25, 50 or 75 mL/m<sup>2</sup>, depending on the location and the situation. Do not apply solution volumes below 2 L/m<sup>2</sup>.

In situations where the soil surface is very dry and conditions are conducive to rapid drying, the area to be treated should be moistened prior to the termiticide application.

It is important to note that when applying a horizontal barrier to the perimeter of a building or structure the chemical barrier is deemed to have a depth of 80 mm. In situations where the solution will not readily wet the soil to the required depth, loosen soil to a depth of 80 mm by 150 mm wide and apply 1.5 L of solution per lineal metre.

**Vertical Barrier Treatments:** To install a vertical barrier use a minimum of 100 L of solution per m<sup>2</sup> of soil. Vertical barriers must be a minimum of 150 mm wide, extend down to 80 mm below the top of the footing and be complete and continuous. Vertical barriers can be installed by trenching and treating the soil as it is backfilled, by soil rodding or by the use of certified reticulation systems, as described in the Australian Standard AS 3660 Series. The preferred method of installing a vertical barrier treatment is either by trenching and treating the soil as it is backfilled or by delivery via a certified reticulation system. When using the soil rodding method to establish a vertical barrier the distance between rod spacings should be as per the following table. To improve soil penetration, the soil should be loosened to a depth of 150mm.

Soil Type	Rod spacing (mm)
Heavy clay	150
Clay loams	200
Loams	250
Sands	300

#### Perimeter Barrier Treatments:

Perimeter barriers consist of horizontal barriers at least 150 mm wide adjoining a vertical barrier of at least 150 mm in width. A perimeter barrier must completely surround all buildings, pipes, piers and service penetrations. In buildings with suspended floors with greater than 400 mm crawl space, perimeter barriers should be installed to surround piers, stumps and service penetrations and completely about all substructure walls.

To ensure provision of a continuous barrier use a minimum of 100 L of solution per m<sup>3</sup> of soil. This equates to a delivery volume of 5 L of solution per linear metre for a 300 mm vertical barrier, or 10 L of solution per linear metre for a 600 mm vertical barrier.

Termites may gain access behind engaged piers against single brick walls unless the soil is treated on both sides of the wall down to the footing.

**Treatment in Conjunction with Physical Barriers:** In situations where the termite protection system is to consist of a combination of both physical and chemical barriers, each certified system must be installed according to the relevant and appropriate product specification and the Australian Standard AS 3660 Series.

**Service Requirements:** Service requirements are to be determined as a result of at least an annual inspection by a licensed Pest Control Operator. More frequent inspections may be required in high risk termite areas.

In determining the need for service, factors such as local termite pressure, breaches of the barrier and termiticide longevity should be considered.

**Subterranean termites are on occasions capable of bridging termite barriers and therefore regular inspections, as detailed in the Australian Standard AS 4349.3, will significantly increase the probability of detection of termite activity before any damage or costly repairs are required.**

Several factors contribute to longevity of the termite treatment and must be considered when evaluating the need for re-treatment. The actual protection period will depend on the termite hazard, climate, soil conditions and rate of termiticide used. Refer to Table A for the expected protection periods provided.

#### INSECTICIDE RESISTANCE WARNING

GROUP	3A	INSECTICIDE
-------	----	-------------

Weed Force Dart 100SC Grub Spider Ant Insecticide is a Group 3A insecticide. Some naturally occurring insect biotypes resistant to Weed Force Dart 100SC Grub Spider Ant Insecticide

and other Group 3A insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if Weed Force Dart 100SC Grub Spider Ant Insecticide or other Group 3A insecticides are used repeatedly. The effectiveness of Weed Force Dart 100SC Grub Spider Ant Insecticide on resistant individuals could be significantly reduced. Since the occurrence of resistant insects is difficult to detect prior to use, Weed Force Pty Ltd accepts no liability for any losses that may result from the failure of Weed Force Dart 100SC Grub Spider Ant Insecticide to control resistant insects. Weed Force Dart 100SC Grub Spider Ant Insecticide may be subject to specific resistance management strategies. For further information contact your local supplier, Weed Force representative or local agricultural department agronomist.

#### **PRECAUTIONS AND RE-ENTRY PERIODS**

DO NOT spray directly on humans, pets or animals. Avoid contact with food, food utensils and preparation surfaces.

Urban pest control: Re-entry: DO NOT allow people and pets to enter treated areas until the spray has dried (normally 3-4 hours) and ventilate buildings before reoccupying. When prior entry is necessary, wear cotton overalls buttoned to the neck and wrist and elbow-length PVC, neoprene or nitrile gloves and chemical resistant footwear. Clothing must be laundered after each day's use.

Re-entry Period for vector control applications: Ventilate indoor treated areas and allow all areas to completely dry (normally 1-2 hours) before re-occupying. When prior entry is necessary, wear cotton overalls buttoned to the neck, wrist and elbow-length PVC, neoprene or nitrile gloves and chemical resistant footwear. Clothing must be laundered after each day's use.

Crops, Ornamentals, Turf: DO NOT allow entry into treated areas until the spray deposits have dried or been watered in after treatment. When prior entry is necessary, wear suitable protective clothing (ie water-proof boots, overalls and gloves). Clothing must be laundered after each day's use.

#### **PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**

Dangerous to fish and aquatic organisms. Do not contaminate dams, rivers, streams, waterways or drains with the product or used container. DO NOT apply to sand, mud, mangrove or aquatic habitats. Drift from treated areas may be hazardous to organisms in adjacent aquatic sites. Extreme caution must be used to avoid aquatic contamination. Avoid spraying flowers that attract and harbour bees.

#### **PROTECTION OF PETS AND LIVESTOCK**

Dangerous to bees. DO NOT spray any plants in flower when bees are foraging. Spray in the night or early morning when bees are not actively foraging. Before spraying remove pets and animals from the areas to be treated. Cover or remove any open food and water containers.

Cover or remove fish ponds, aquariums etc. before spraying. Do not allow re-entry until spray has dried.

#### **STORAGE AND DISPOSAL**

Store in closed original containers, in a cool, well ventilated area away from children, animals, food and feedstuffs. Do not store for prolonged periods in direct sunlight. In case of spillage, confine and absorb spilled product with absorbent material such as sand, clay or cat litter. Dispose of waste as indicated below or according to the Australian Standard AS 2507 - Storage and Handling of Pesticides. Do NOT allow spilled product to enter sewers, drains, creeks or any other waterways.

Triple rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulation. Do not burn empty containers or product.

For refillable containers empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

#### **SAFETY DIRECTIONS**

Poisonous if swallowed. May irritate the eyes and skin. Repeated exposure may cause allergic disorders. Avoid contact with eyes and skin.

For termite control in buildings and structures: When opening the container, preparing spray and using prepared spray, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow length PVC or nitrile gloves. After each day's use, wash gloves and contaminated clothing.

For handheld application: When opening the container and preparing spray, wear cotton overalls buttoned to the neck and wrist, washable hat and elbow length PVC or nitrile gloves. When using prepared spray wear protective waterproof clothing, elbow length PVC or nitrile gloves and water resistant footwear.

After each day's use, wash gloves and contaminated clothing. Wash hands after use.

#### **FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126; New Zealand 0800 764 766.



### **SAFETY DATA SHEET (SDS)**

The SDS contains important information relating to health and safety. This product should be used in conjunction with information on the SDS and users are advised to read the SDS before using this product. The most up to date SDS can be obtained from [www.weedforce.com.au](http://www.weedforce.com.au)

### **CONDITIONS OF SALE**

To the extent permitted by law, all conditions and warranties and statutory or other rights or action which buyer or any other user may have against Weed Force Pty Ltd ("Weed Force") or Seller are hereby excluded. Weed Force hereby gives Notice to Buyer and other users that it will not accept responsibility for any direct, indirect or consequential loss however arising (including negligence) from the sale, supply, use or application of this product or reliance on product information or advice provided by Weed Force or on its behalf. No warranty expressed or implied is given by Weed Force regarding its suitability or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not. Weed Force's liability shall, in all circumstances, be limited to replacement of the product or a refund of the purchase price paid therefor.

**WEEDFORCE** Pty Ltd **ACN 602 207 152**   
2072 Logan Rd Mt Gravatt QLD 4122  
P: 0488 997 923 [www.weedforce.com.au](http://www.weedforce.com.au)

APVMA Approval No:  
92587 / 135972

® Registered Trademark of Weed Force Pty Ltd  
© Weed Force Pty Ltd 2022